

Assessment of microbial communities associated with fermentative–methanogenic biodegradation of aromatic hydrocarbons in groundwater contaminated with a biodiesel blend (B20)


Débora Toledo Ramos, Márcio Luís Busi da Silva, Carlos Wolfgang Nossa, Pedro J. J. Alvarez & Henry Xavier Corseuil

Biodegradation

ISSN 0923-9820

Biodegradation

DOI 10.1007/s10532-014-9691-4



Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media Dordrecht. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Assessment of microbial communities associated with fermentative–methanogenic biodegradation of aromatic hydrocarbons in groundwater contaminated with a biodiesel blend (B20)

Débora Toledo Ramos · Márcio
Luís Busi da Silva · Carlos Wolfgang Nossa ·
Pedro J. J. Alvarez · Henry Xavier Corseuil

Received: 5 December 2013 / Accepted: 9 April 2014
© Springer Science+Business Media Dordrecht 2014

Abstract A controlled field experiment was conducted to assess the potential for fermentative–methanogenic biostimulation (by ammonium–acetate injection) to enhance biodegradation of benzene, toluene, ethylbenzene and xylenes (BTEX) as well as polycyclic aromatic hydrocarbons (PAHs) in groundwater contaminated with biodiesel B20 (20:80 v/v soybean biodiesel and diesel). Changes in microbial community structure were assessed by pyrosequencing 16S rRNA analyses. BTEX and PAH removal began 0.7 year following the release,

concomitantly with the increase in the relative abundance of *Desulfitobacterium* and *Geobacter* spp. (from 5 to 52.7 % and 15.8 to 37.3 % of total Bacteria 16S rRNA, respectively), which are known to anaerobically degrade hydrocarbons. The accumulation of anaerobic metabolites acetate and hydrogen that could hinder the thermodynamic feasibility of BTEX and PAH biotransformations under fermentative/methanogenic conditions was apparently alleviated by the growing predominance of *Methanosarcina*. This suggests the importance of microbial population shifts that enrich microorganisms capable of interacting syntrophically to enhance the feasibility of fermentative–methanogenic bioremediation of biodiesel blend releases.

D. T. Ramos (✉) · H. X. Corseuil
Department of Sanitary and Environmental Engineering,
Federal University of Santa Catarina, Florianópolis, SC,
Brazil
e-mail: debora.toledo@uol.com.br

H. X. Corseuil
e-mail: henry.corseuil@ufsc.br

M. L. B. da Silva
EMBRAPA, BR 153 km 110, P.O. Box 21, Concórdia,
SC 89700-000, Brazil
e-mail: marcio.busi@embrapa.br

C. W. Nossa
Department of Ecology and Evolutionary Biology,
Rice University, Houston, TX, USA
e-mail: carlosn@genebygene.com

P. J. J. Alvarez
Department of Civil and Environmental Engineering,
Rice University, Houston, TX, USA
e-mail: alvarez@rice.edu

Keywords Biodegradation · Biodiesel · BTEX ·
PAH · Pyrosequencing · Syntrophy

Introduction

The use of biodiesel blends as an alternative renewable transportation fuel is increasing worldwide to alleviate dependence on fossil fuels and to minimize atmospheric emissions and greenhouse effects. The increasing biodiesel demand can, however, increase the probability of groundwater contamination as result of accidental and incidental spills during its production, transportation and storage. Although biodiesel is commonly referred to as a harmless and readily

Table 1 Main reactions involved in biodiesel, BTEX and PAH degradation and ΔG° values (kJ mol^{-1})

Reactions involved in linoleic acid (1, 2, 3 and 4), benzene (5, 6, 7 and 8) and naphthalene (9, 10, 11 and 12) degradation	ΔG° reaction ^a (kJ mol^{-1})
Linoleic acid (C _{18:2})	
(1) C ₁₈ H ₃₁ O ₂ ⁻ + H ⁺ + 16H ₂ O → 9CH ₃ COO ⁻ + 9H ⁺ + 14H ₂	+272.33
(2) 9CH ₃ COO ⁻ + 9H ⁺ + 18H ₂ O → 36H ₂ + 18CO ₂	+854.26
(3) 50H ₂ + 12,5CO ₂ → 12,5CH ₄ + 25H ₂ O	-1634.34
(4) Sum (1) + (2) + (3): C ₁₈ H ₃₁ O ₂ ⁻ + H ⁺ + 9H ₂ O → 12,5CH ₄ + 5,5CO ₂	-507.75
Benzene	
(5) C ₆ H ₆ + 6H ₂ O → 3CH ₃ COO ⁻ + 3H ⁺ + 3H ₂	+70.73
(6) 3CH ₃ COO ⁻ + 3H ⁺ + 6H ₂ O → 12H ₂ + 6CO ₂	+284.754
(7) 15H ₂ + 3,75CO ₂ → 3,75CH ₄ + 7,5H ₂ O	-490.30
(8) Sum (5) + (6) + (7): C ₆ H ₆ + 4,5 H ₂ O → 3,75CH ₄ + 2,25CO ₂	-134.82
Naphthalene	
(9) C ₁₀ H ₈ + 10H ₂ O → 5CH ₃ COO ⁻ + 5H ⁺ + 4H ₂	+101.12
(10) 5CH ₃ COO ⁻ + 5H ⁺ + 10H ₂ O → 10CO ₂ + 20H ₂	+474.59
(11) 24H ₂ + 6CO ₂ → 6CH ₄ + 12H ₂ O	-784.48
(12) Sum (9) + (10) + (11): C ₁₀ H ₈ + 8H ₂ O → 6CH ₄ + 4CO ₂	-291.75

$\Delta G_f^{\circ}(\text{aq})$ values of linoleic acid and naphthalene were obtained in Lalman (2000) and Doling et al. (2009), respectively. All other compounds ΔG_f° (aq and g (for H₂)) values were obtained in Thauer et al. (1977)

^a Standard Gibbs energies were calculated under standard conditions (1 M solute concentration, pH = 7, T = 298 K and gas partial pressure of 1 atm)

biodegradable biofuel (Zhang et al. 1998), it is usually blended with petroleum diesel fuel that contains priority pollutants such as benzene, toluene, ethylbenzene and xylenes (BTEX) and polycyclic aromatic hydrocarbons (PAH). These hydrocarbons include carcinogenic compounds (e.g., benzene and benzo[*a*]pyrene) that are generally monitored to determine the need for corrective remedial action.

The high biochemical oxygen demand exerted by indigenous microorganisms during biodiesel biodegradation rapidly drives impacted aquifers towards methanogenic conditions. This phenomenon is particularly noticeable at the source zone region where higher concentration of organic compounds stimulates the consumption of terminal electron acceptors. Under methanogenic conditions where the energetic yield is close to the minimum needed for microbial sustenance ($\approx -20 \text{ kJ mol}^{-1}$ required for ATP formation) (Schink 1997), bioremediation is usually accomplished by syntrophic microorganisms (Morris et al. 2013). Syntrophic anaerobes can play a critical role in the biodegradation of long-chain fatty acids (Sousa et al. 2009), BTEX (Rakoczy et al. 2011) and PAH (Berdugo-Clavijo et al. 2012), since initial fermentative/

methanogenic biotransformations are thermodynamically unfeasible (endergonic) (Table 1, reactions 1, 5 and 9) without consumption of degradation byproducts by commensal microorganisms. Long-chain fatty acids (represented by linoleic acid) derived from biodiesel esters hydrolysis can be further oxidized to acetate and hydrogen via β -oxidation (Sousa et al. 2009). Although this reaction is thermodynamically unfeasible (Table 1, reaction 1), it might proceed if syntrophic microorganisms consume metabolites that can impose thermodynamic constraints (reactions 2 and 3) thus making the overall reaction exergonic (reaction 4). Similarly, fermentative/methanogenic BTEX and PAH biodegradation (represented by benzene and naphthalene, respectively) is plausible (reactions 8 and 12) when syntrophic microorganisms consume the metabolites (reactions 6, 7, 10 and 11).

We previously demonstrated that anaerobic biostimulation by the addition of ammonium acetate into B20 contaminated groundwater induced fermentative-methanogenic conditions that enhanced BTEX removal (Ramos et al. 2013). This enhancement in BTEX anaerobic biodegradation was hypothesized to occur due to the proliferation of putative hydrocarbon

degraders thriving syntrophically with methanogenic archaea. Nonetheless, the microbial community structure was not characterized to identify potential syntrophic associations. Therefore, in this work, microbial 16S rRNA pyrosequencing analyses were conducted to assess temporal changes in microbial community structure during anaerobic biostimulation of groundwater contaminated with a biodiesel blend. Emphasis was placed on studying microbial populations putatively associated with aromatic hydrocarbon biodegradation. This information advances our current understanding of fermentative–methanogenic bioprocesses by identifying dominant microorganisms during anaerobic bioremediation of B20 releases.

Materials and methods

Controlled release field experiment

A controlled release field experiment was monitored over 2 years to investigate whether anaerobic biostimulation could enhance BTEX biodegradation under fermentative–methanogenic conditions. Detailed information on this experiment set up and groundwater monitoring was previously described (Ramos et al. 2013). Briefly, a source zone was established by releasing 100L of B20 (20 % v/v soybean biodiesel and 80 % v/v diesel) into an area of 1 m² × 1.6 m deep down to the water table. Fermentative–methanogenic biostimulation was performed by weekly injection of ammonium acetate (300 mg L⁻¹) into 5 wells installed 1.5 m upstream of the source zone.

Chemical analyses

Groundwater was monitored at the source zone since biodiesel blend releases are not readily miscible in groundwater and behave as a fixed, decaying, yet long-lived source with relatively small region of influence compared to soluble biofuels such as ethanol (Corseuil et al. 2011). Groundwater pH, redox potential, dissolved oxygen, nitrite, nitrate, sulfate, sulfide, ferrous iron, acetate, BTEX and methane were monitored over time (Ramos et al. 2013).

PAH were extracted from groundwater using solid phase SPE cartridges, according to EPA method 525.2, and measured by gas chromatography (HP model 6890 II with a flame ionization detector (FID)

and HP-5 capillary column). Detection limits were (in parenthesis): naphthalene (7 µg L⁻¹), methylnaphthalene (5 µg L⁻¹), dimethylnaphthalene (7 µg L⁻¹), acenaphthylene (8 µg L⁻¹), acenaphthene (8 µg L⁻¹), fluorene (8 µg L⁻¹), phenanthrene (9 µg L⁻¹), anthracene (9 µg L⁻¹), fluoranthene (10 µg L⁻¹), pyrene (9 µg L⁻¹), benzo[*a*]anthracene (9 µg L⁻¹), chrysene (10 µg L⁻¹), dibenzo[*a,h*]anthracene (12 µg L⁻¹), benzo[*b*]fluoranthene (12 µg L⁻¹), benzo[*k*]fluoranthene (31 µg L⁻¹), benzo[*a*]pyrene (36 µg L⁻¹), indeno[1,2,3-*cd*]pyrene (28 µg L⁻¹) and benzo[*g,h,i*]pyrene (11 µg L⁻¹).

Dissolved hydrogen analyses were conducted 2.4 years following the B20 release using in situ passive samplers. These samplers were deployed for 5 days in groundwater, according to the methods developed by Spalding and Watson 2006 and Spalding and Watson 2008. Hydrogen was measured by a gas chromatography (model UC-13 Construmaq equipped with a 30-ft long × 1/8 in stainless-steel column packed with 100/120 mesh Haye SepD solid phase and thermal conductivity detector). The detection limit was 10⁻⁹ M.

Microbial analysis

16S rRNA pyrosequencing analyses were conducted to identify, characterize and assess temporal changes in microbial community structure during anaerobic biostimulation of groundwater contaminated with a biodiesel blend. DNA samples were extracted from groundwater before the B20 release as background information on indigenous microorganisms and then after 0.3, 0.7, 1.0, 1.4 and 1.6 years following the fuel release.

To amplify the 16S rRNA region two primer pairs were used: one for Bacterial 16S and one for Archaeal 16S. Bacteria 16S rRNA was amplified within hyper variable region 3 and 4 (bases 347-803) using primers designed specifically for 454 sequencing (which included 454 adapter sequence, 10 base nucleotide barcode, and target primer). Target primers were: forward 5'-GGAGGCAGCAGTRRGGAAT-3', reverse 5'-CTACCRGGGTATCTAATCC-3' (Nossa et al. 2010). Archaea 16S was amplified between bases 571–1,204. Target primers were: forward 5'-GCYTAAAGSRICCGTAGC-3', reverse 5'-TTMGGGGCATRCIKACCT-3' (Gantner et al. 2011). PCR was carried out using Qiagen Top Taq

Table 2 Metabolic characteristics of the bacteria genera found in groundwater source-zone

Bacteria	Class	Respiration mode	Substrates	References
<i>Janthinobacterium</i>	β -Proteobacteria	Aerobic	PAH	Bodour et al. (2003), Lima et al. (2012)
<i>Ralstonia</i>	β -Proteobacteria	Aerobic or denitrifier	Fatty acids, alcohols, aromatic compounds	Cramm (2009), Bodour et al. (2003)
<i>Burkholderia</i>	β -Proteobacteria	Facultative	Methyl esters, acetate, alkanes, aromatic compounds and chlorinated compounds	Philippe et al. (2001), O' Sullivan and Mahenthiralingam (2005), Belova et al. (2006)
<i>Geobacter</i>	δ -Proteobacteria	Iron-reducer, fermentative	Short-chain fatty acids, alcohols and monoaromatic compounds	Lovley et al. (1993), Cord-Ruwisch et al. (1998)
<i>Geobacillus</i>	Bacilli	Facultative	Organic acids, alcohols, aromatic compounds and carbohydrates	Nazina et al. (2001)
<i>Staphylococcus</i>	Bacilli	Facultative	Sugars	Kloos (1980)
<i>Desulfotobacterium</i>	Clostridia	Sulfate-, thiosulfate-, sulfite-, or iron-reducer, fermentative	Halogenated organic compounds, aromatic compounds, organic acids	Villemur et al. (2006), Kunapuli et al. (2010)
<i>Gp 1, 3, 13, 23</i>	Acidobacteria	Facultative	Sugars, amino acids, alcohols, organic acids	Ward et al. (2009), Geissler et al. (2009)
<i>Nitrospira</i>	Gp 1, 3, 13, 23	Aerobic or nitrite oxidation	Nitrogenated compounds	Blackbourne et al. (2007)
<i>Pseudomonas</i>	γ -Proteobacteria	Facultative denitrifier	Aromatic compounds	Shim et al. (2005), Bruce et al. (2010), Lima et al. (2012)
<i>Desulfovibrio</i>	δ -Proteobacteria	Sulfate-, thiosulfate- or sulfite reducer, fermentative	Long-chain fatty acids, aromatic compounds, organic acids, alcohol	Allen et al. (2008), Sousa et al. (2007), Heidelberg et al. (2004), Bryant et al. (1977)
<i>Desulfosporosinus</i>	Clostridia	Sulfate-, thiosulfate-, sulfite-, or arsenate-reducer	Sugars, alcohols, monoaromatic compounds	Robertson et al. (2001), Liu et al. (2004), Ramamoorthy et al. (2006)
<i>Clostridium III</i>	Clostridia	Fermentative	Long-chain fatty acids, organic acids	Chauhan and Ogram (2006), Hatamoto et al. (2007)
<i>Geothrix</i>	Holophagae	Iron-reducer or fermentative	Organic acids	Coates et al. (1999), Nevin and Lovley (2002)
<i>Anaeromyxobacter</i>	δ -Proteobacteria	Denitrifier or iron-reducer	Chlorinated organic compounds, organic acids	Treude et al. (2003), Sanford et al. (2002), He and Sanford (2004)
<i>Desulfomonile</i>	δ -Proteobacteria	Sulfate-reducer	Chlorinated organic compounds	Dolfing and Tiedje (1991)
<i>Petotomaculum</i>	Clostridia	Fermentative	Organic acids, alcohols	Imachi et al. (2002)
<i>Holophaga</i>	Holophagae	Fermentative	Short-chain fatty acids, methylated aromatic compounds	Liesack et al. (1994), Chauhan and Ogram (2006)

Table 2 continued

Bacteria	Class	Respiration mode	Substrates	References
<i>Peptococcaceae</i>	Clostridia	Denitrifier, sulfate- or iron-reducer	Aromatic compounds	van der Zaan et al. (2012)
<i>Microvirgula</i>	β -Proteobacteria	Aerobic or denitrifier	Organic acids and alcohols	Patureau et al. (1998)
<i>Curvibacter</i>	β -Proteobacteria	Aerobic	Sugars, amino acids	Ding and Yokota (2010)
<i>Neusikia</i>	γ -Proteobacteria	Aerobic	Carbohydrates	Leandro et al. (2012)
<i>Desulfovirga</i>	δ -Proteobacteria	Sulfate-, thiosulfate-, or sulfite-reducer	Short-chain fatty acids, alcohols	Tanaka et al. (2000)

PCR kit as per manufacturer's specifications (Qiagen, Carlsbad, California). Thermal cycling was carried out at 94 °C for 3 min, followed by 30 cycles of (94 °C for 30 s, 55 °C for 30 s, 72 °C for 60 s) followed by 10 min at 72 °C and hold at 4 °C. PCR products were run on an agarose gel and the amplicons verified for size against DNA ladder and further excised for purification. Gel bands were purified using the Qiagen Qiaquick gel extraction kit as per manufacturer's specifications. Purified DNA was eluted with TE buffer, and quantified using Picogreen fluorescent DNA quantitation kit. 16S amplicons from different samples were pooled in equimolar concentrations prior to emulsion PCR and sequencing using Roche 454 Jr. Image data was processed using the Amplicon processing pipeline in the 454 Jr. (Roche, Branford, CT) suite of processing software included with the sequencer.

Raw sequencing data was initially processed using the Ribosomal Database Project (RDP) Pyrosequencing pipeline (<http://pyro.cme.msu.edu>) (Cole et al. 2009). Raw sequences were first sorted by barcode, and fusion primers were removed. Quality filter removed sequences with lengths less than 150 bases or having more than one ambiguity (*N*) or those with more than two changes in forward or reverse primers. Filtered, trimmed sequence data was then classified using RDP classifier at the genus level.

Results and discussion

Eight dominant phylogenetic classes constituted the groundwater microbial community: β -Proteobacteria, γ -Proteobacteria, δ -Proteobacteria, Bacilli, Clostridia, Acidobacteria, Nitrospira and Holophagae (Table 2). Prior to B20 fuel release, microaerophilic conditions prevailed with groundwater dissolved oxygen concentration of 0.6 mg L⁻¹ and redox potential of +104 mV (Table 3). Such microaerophilic conditions likely favored the growth of aerobic facultative hydrocarbon degrading bacteria, including some *Janthinobacterium*, *Ralstonia*, *Pseudomonas* and *Geobacillus* species (Fig. 1). After 0.3 year following the fuel release, *Burkholderia* was the main bacterial genus detected. This bacterium is generally capable of degrading a variety of organic compounds (Philippe et al. 2001; Belova et al. 2006) (Table 2). Nonetheless, negligible BTEX and PAH removal was observed

Table 3 Temporal shifts in groundwater geochemical conditions at the source zone (2 m below ground surface)

Time (years)	Temperature (°C)	pH	Redox potential (mV)	DO (mg L ⁻¹)	NO ₃ ⁻ (mg L ⁻¹)	NO ₂ ⁻ (mg L ⁻¹)	Fe II (mg L ⁻¹)	SO ₄ ²⁻ (mg L ⁻¹)	S ²⁻ (mg L ⁻¹)	Acetate (mg L ⁻¹)	CH ₄ (mg L ⁻¹)
0	21.8	4.3	+104	0.62	16.8	<0.1	6.1	2.2	0.01	<0.1	<0.005
0.3	21.1	3.9	+31	0.35	<0.1	<0.1	174.3	1.1	0.04	131	1.5
0.7	25.8	4.5	-36	0.77	<0.1	<0.1	13.9	0.8	0.01	5.3	17.5
1.0	27.2	4.8	-131	0.40	<0.1	<0.1	4.1	5.2	0.02	4.2	8.5
1.4	23.1	4.6	-77	0.26	<0.1	<0.1	4.5	0.9	0.01	16.8	3.7
2.0	26.6	4.9	-140	0.08	<0.1	<0.1	6.5	0.5	0.03	8.9	7.2
2.4	24.6	4.5	-73	0.4	<0.1	<0.1	9.2	0.5	0.30	3.9	6.2

during the time frame in which *Burkholderia* was predominant (Figs. 1, 2). Therefore, it is plausible that the proliferation of *Burkholderia* was primarily associated with the consumption of acetate or biodiesel methyl-esters.

Noticeable BTEX removal occurred at 0.7 year, considerably faster than under natural attenuation conditions (with a noticeable onset of BTEX biodegradation after 2.5 years) (Ramos et al. 2013). BTEX removal was accompanied by an increase in the relative abundance of *Desulfitobacterium* and *Geobacter* (from 5 to 52.7 and 15.8 to 37.3 % of total Bacteria 16S rRNA, respectively) (Fig. 1). These bacteria have been widely reported to participate in the anaerobic biodegradation of aromatic hydrocarbons under iron-reducing conditions (Lovley et al. 1993; Coates et al. 1995; Kunapuli et al. 2010), which is consistent with the observed accumulation of iron (II) (Table 3). *Geobacter* and *Desulfitobacterium* remained abundant even after the establishment of fermentative conditions (Fig. 2), suggesting that these organisms may have relied on fermentative metabolism as well (Cord-Ruwisch et al. 1998; Kunapuli et al. 2010). It should be noted that bacteria belonging to δ -Proteobacteria (iron and sulfate-reducing genera including *Geobacter*, *Pelobacter*, *Desulfovibrio*, *Desulfomicrobium*, *Desulfuromusa* and *Desulfuromonas*) were below detection limit (10^2 gene copies g⁻¹) in the experimental control plot not biostimulated (Ramos et al. 2013).

The predominance of methanogenic archaeal communities coincided with the establishment of methanogenic conditions in groundwater (i.e., 17.5 mg CH₄ L⁻¹ measured at 0.7 year following the fuel release (Table 3)). Archaea community analysis revealed the presence of 7 different genera: *Thermoprotei*, *Methanosarcinaceae*, *Thermogymnomonas*, *Methanosaeta*, *Methanosarcina*, *Methanospirillum* and *Methanoregula* (Fig. 3). *Thermoprotei*, which can thrive using iron (III) as a terminal electron acceptor (Slobodkin 2005; Wagner and Wiegel 2008) were first detected after 0.3 year following the release (Fig. 1) and concomitantly with the prevalence of iron-reducing conditions (Fig. 2; Table 3). Both *Methanosaeta* and *Methanosarcina* are commonly found in acetate-rich methanogenic environments (Liu and Whitman 2008). The increased abundance of *Methanosarcina* compared to aceticlastic *Methanosaeta* (Liu and Whitman 2008) (Fig. 3) could be

Fig. 1 Temporal changes in 16S rRNA relative abundance (%) of bacteria communities in groundwater samples from the B20 source zone





Fig. 2 **a** Iron (II), BTEX and total PAH concentrations and associated changes in the abundance of the putative aromatic hydrocarbon degraders *Geobacter* and *Desulfitobacterium*, expressed as a % of the total Bacteria 16S rRNA. **b** Acetate and methane profiles. The dashed horizontal line is the methane solubility limit (22 mg L⁻¹ at 24 °C and 1 atm). Arrows indicate the start and the end of biostimulation with ammonium acetate

attributable to their faster growth rates when acetate is not limiting (Smith and Ingram-Smith 2007) and versatile metabolism (e.g., hydrogenotrophic, acetate-fermenting or methylotrophic while *Methanosaeta* is exclusively acetate-fermenting) (Galagan et al. 2002; Liu and Whitman 2008).

One year after the release, *Methanoregula* and *Methanospirillum* were detected in groundwater (Fig. 3). The presence of these hydrogenotrophic microorganisms (Worm et al. 2011; Bräuer et al. 2011) suggests their role as H₂ consumers that could alleviate the thermodynamic constraints caused by H₂ accumulation (Table 1, reactions 3, 7 and 11).

Syntrophic cooperation between aromatic hydrocarbons degraders and methanogens

Syntrophic relationships allow microorganisms to thrive in energetically constrained systems (e.g., methanogenic environments). To discern whether a biological process meets the requirements for being considered syntrophy, one may consider the reactions involved and discern whether they are feasible without microbial cooperation (Morris et al. 2013). Table 1 summarizes the need for syntrophic cooperation to enhance the thermodynamic feasibility of biodiesel, BTEX and PAH biodegradation.

Fig. 3 Temporal changes in 16S rRNA relative abundance (%) of Archaea communities in groundwater samples from the B20 source zone





Fig. 4 Thermodynamic feasibility of benzene (0.9 mg L⁻¹) (a) and naphthalene (0.16 mg L⁻¹) (b) fermentation to acetate and hydrogen (Table 1, reactions 5 and 9) for typical H₂ concentrations of 10⁻⁷ to 10⁻⁹ M (grey diagonal lines), dissolved H₂ concentration detected in groundwater at 2.4 years (black diagonal lines) and pH = 4.5. Grey vertical lines represent acetate concentration at 0.3 (131 mg L⁻¹), 0.7 (5 mg L⁻¹) and 1.0 year (8.9 mg L⁻¹) after the release. Black vertical lines correspond to acetate concentration at 2.4 years (3.9 mg L⁻¹)



Thermodynamic calculations considered the highest contaminant concentrations (0.9 mg L⁻¹ for benzene and 0.16 mg L⁻¹ for naphthalene) and typical dissolved H₂ concentrations found in anaerobic aquifers (10⁻⁷ to 10⁻⁹ M) (Heimann et al. 2009), as well as dissolved H₂ concentrations detected in groundwater 2.4 years after the release. As shown in Fig. 4a, the presence of 131 mg L⁻¹ of acetate (at 0.3 year) hindered fermentative/methanogenic BTEX biodegradation (represented by benzene) for dissolved H₂ concentrations $\geq 0.2 \times 10^{-8}$ M. With a decrease in acetate concentrations from 131 to 5 mg L⁻¹ (measured at 0.7 year) benzene biodegradation became thermodynamically feasible. These predictions were in agreement with the observed in situ BTEX removal pattern shown in Fig. 2. Biodegradation of PAH (represented by naphthalene, 0.16 mg L⁻¹) was thermodynamically feasible for almost all H₂ concentrations except 10⁻⁷ M (at 1.0 year) (Fig. 4b).

Conclusions

This assessment of microbial community structure over time advances our understanding of the syntrophic metabolic niches that evolve during fermentative–methanogenic bioremediation of aromatic hydrocarbons. Microbial population shifts were consistent with observed geochemical changes and dominance shifted towards the putative anaerobic hydrocarbon degraders *Geobacter* and *Desulfitobacterium*. The accumulation of anaerobic metabolites acetate and H₂ that could thermodynamically hinder BTEX and PAH biodegradation was likely alleviated by the proliferation of *Geobacter*, *Desulfitobacterium* and hydrogenotrophic or acetoclastic *Methanosarcina*. Overall, this study suggests that fermentative–methanogenic biostimulation can promote favorable microbial population shifts that enhance the natural attenuation of B20 blend releases.

Acknowledgments The authors thank PETROBRAS (Petróleo Brasileiro S/A) for the research financial support as well as CAPES (Coordination for the Improvement of Higher Level Education Personnel) and CNPq (National Council for Scientific and Technological Development) for providing scholarships.

References

- Allen TD, Kraus PF, Lawson PA, Drake GR, Balkwill DL, Tanner RS (2008) *Desulfovibrio carbinoliphilus* sp., a benzyl alcohol-oxidizing, sulfate-reducing bacterium isolated from a gas condensate-contaminated aquifer. *Int J Syst Evol Microbiol* 58(6):1313–1317
- Belova SE, Pankratov TA, Dedysh SN (2006) Bacteria of the genus *Burkholderia* as a typical component of the microbial community of *Sphagnum* peat bogs. *Microbiology* 75(1):90–96
- Berdugo-Clavijo C, Dong X, Soh J, Sensen CW, Gieg LM (2012) Methanogenic biodegradation of two-ringed polycyclic aromatic hydrocarbons. *FEMS Microbiol Ecol* 81(1):124–133
- Blackbourne R, Vadivelu VM, Yuan Z, Keller J (2007) Kinetic characterisation of an enriched *Nitrospira* culture with comparison to *Nitrobacter*. *Water Res* 41(14):3033–3042
- Bodour AA, Wang JM, Brusseau ML, Maier RM (2003) Temporal change in culturable phenanthrene degraders in response to long-term exposure to phenanthrene in a soil column system. *Environ Microbiol* 5(10):888–895
- Bräuer SL, Cadillo-Quiroz H, Ward RJ, Yavitt JB, Zinder SH (2011) *Methanoregula boonei* gen. nov., sp. nov., an acidiphilic methanogen isolated from an acidic peat bog. *Int J Syst Evol Microbiol* 61(1):45–52
- Bruce T, Martinez IB, Neto OM, Vicente ACP, Kruger RH, Thompson FL (2010) Bacterial community diversity in the Brazilian Atlantic forest soils. *Soil Microbiol* 60(4):840–849
- Bryant MP, Campbell LL, Reddy CA, Crabill MR (1977) Growth of *Desulfovibrio* in lactate or ethanol media low in sulfate in association with H₂-utilizing methanogenic bacteria. *Appl Environ Microbiol* 33(5):1162–1169
- Chauhan A, Ogram A (2006) Fatty acid-oxidizing consortia along a nutrient gradient in the Florida Everglades. *Appl Environ Microbiol* 72(4):2400–2406
- Coates JD, Lonergan DJ, Phillips EJP, Jenter A, Lovley DR (1995) *Desulfuromonas palmitatis* sp., nov. A marine dissimilatory Fe(III) reducer that can oxidize long chain fatty acids. *Arch Microbiol* 164(6):406–413
- Coates JD, Ellis DJ, Gaw CV, Lovley DL (1999) *Geothrix fermentans* gen. nov., sp. nov., a novel Fe(III)-reducing bacterium from a hydrocarbon-contaminated aquifer. *Int J Syst Evol Microbiol* 49(4):1615–1622
- Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ, Kulam-Syed-Mohideen AS, McGarrell DM, Marsh T, Garrity GM, Tiedje JM (2009) The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res* 37:141–145
- Cord-Ruwisch R, Lovley DR, Schink B (1998) Growth of *Geobacter sulfurreducens* with acetate in syntrophic cooperation with hydrogen-oxidizing anaerobic partners. *Appl Environ Microbiol* 64(6):2232–2236
- Corseuil HX, Monier AL, Gomes APN, Chiaranda HS, Rosario M, Alvarez PJJ (2011) Biodegradation of soybean and castor oil biodiesel: implications on the natural attenuation of monoaromatic hydrocarbons in groundwater. *Groundw Monit Remediat* 31(3):111–118
- Cramm R (2009) Genomic view of energy metabolism in *Ralstonia eutropha* H16. *J Mol Microbiol Biotechnol* 16(1–2):38–52
- Ding L, Yokota A (2010) *Curvibacter fontana* sp. nov., a microaerobic bacterium isolated from well water. *J Gen Appl Microbiol* 56(3):267–271

- Dolfing J, Tiedje JM (1991) Acetate as a source of reducing equivalents in the reductive dechlorination of 2,5-dichlorobenzoate. *Arch Microbiol* 156(5):356–361
- Dolfing J, Xu A, Gray ND, Larter SR, Head IM (2009) The thermodynamic landscape of methanogenic PAH degradation. *Microb Biotechnol* 2(5):566–574
- Galagan JE, Nusbaum C, Roy A, Endrizzi MG, MacDonalds P, Fitzhugh W, Calvo S, Engels R, Smirnov S, Atnoor D, Brown A, Allen N, Naylor J, Stange-Thomann N, Dearlano K, Johnson R, Linton L, Mcewan P, Mckernan K, Talamas J, Tirrell A, Ye WJ, Zimmer A, Barber RD, Cann I, Graham DE, Grahame DA, Guss AM, Hedderich R, Ingram-Smith C, Kuettner HC, Krzycki JA, Leigh JA, Li WX, Liu JF, Mukhopadhyay B, Reeve JN, Smith K, Springer TA, Umayam LA, White O, White RH, De Macario EC, Ferry JG, Jarrell KF, Jing H, Macario AJL, Paulsen I, Pritchett M, Sowers KR, Swanson RV, Zinder SH, Lander E, Metcalf WW, Birren B (2002) The genome of *M. acetivorans* reveals extensive metabolic and physiological diversity. *Genome Res* 12(4):532–542
- Gantner S, Andersson AF, Alonso-Sáez L, Bertilsson S (2011) Novel primers for 16S rRNA-based archaeal community analyses in environmental samples. *J Microbiol Methods* 84(1):12–18
- Geissler A, Merroum M, Geipel G, Reuther H, Selenska-Pobell S (2009) Biogeochemical changes induced in uranium mining waste pile samples by uranyl nitrate treatments under anaerobic conditions. *Geobiology* 7(3):282–294
- Hatamono M, Imachi H, Yashiro Y, Ohashi A, Harada H (2007) Diversity of anaerobic microorganisms involved in long-chain fatty acid degradation in methanogenic sludges as revealed by RNA-based stable isotope probing. *Appl Environ Microbiol* 73(13):4119–4127
- He Q, Sanford RA (2004) Acetate threshold concentrations suggest varying energy requirements during anaerobic respiration by *Anaeromyxobacter dehalogenans*. *Appl Environ Microbiol* 70(11):6940–6943
- Heidelberg JF, Seshadri R, Haveman SA, Hemme CL, Paulsen IT, Kolonay JF, Eisen JA, Ward N, Methe B, Brinkac LM, Daugherty SC, Deboy RT, Dodson RJ, Durkin AS, Madupu R, Nelson WC, Sullivan SA, Fouts D, Haft DH, Selengut J, Peterson JD, Davidsen TM, Zafar N, Zhou L, Radune D, Dimitrov G, Hance M, Tran K, Khouri H, Gill J, Utterback TR, Feldblyum TV, Wall JD, Voordouw G, Fraser CM (2004) The genome sequence of the anaerobic, sulfate-reducing bacterium *Desulfovibrio vulgaris*. *Nat Biotechnol* 22(5):554–559
- Heimann A, Jakobsen R, Blodau C (2009) Energetics constraints on H₂-dependent terminal electron accepting processes in anoxic environments: a review of observations and model approaches. *Environ Sci Technol* 44(1):24–33
- Imachi H, Sekiguchi Y, Kamagata Y, Hanada S, Ohashi A, Harada H (2002) *Pelotomaculum thermopropionicum* gen. nov., sp. nov., an anaerobic, thermophilic, syntrophic propionate-oxidizing bacterium. *Int J Syst Evol Microbiol* 52(5):1729–1735
- Kloos WE (1980) Natural populations of the genus *Staphylococcus*. *Annu Rev Microbiol* 34:559–592
- Kunapuli U, Jahn MK, Lueders T, Geyer R, Hermann JH, Meckenstock RU (2010) *Desulfitobacterium aromaticivorans* sp. nov. and *Gebacter toluenoxydans* sp. nov., iron-reducing bacteria capable of anaerobic degradation of monoaromatic hydrocarbons. *Int J Syst Evol Microbiol* 60(3):686–695
- Lalman JD (2000) Anaerobic degradation of linoleic (C18:2), oleic (C18:1) and stearic (C18:0) acids and their inhibitory effects on acidogens, acetogens and methanogens. Thesis. University of Toronto
- Leandro T, França L, Nobre MF, Schumann P, Rosselló-Móra R, Costa MS (2012) *Nevskia aquatilis* sp. nov. and *Nevskia persephonica* sp. nov., isolated from a mineral water aquifer and the emended description of the genus *Nevskia*. *Syst Appl Microbiol* 35(5):297–301
- Liesack W, Bak F, Kreft JU, Stackebrandt E (1994) *Holophaga foetida* gen. nov. sp. nov., a new homoacetogenic bacterium degrading methoxylated aromatic compounds. *Arch Microbiol* 162(1–2):85–90
- Lima G, Parker B, Meyer J (2012) Dechlorinating microorganisms in a sedimentary rock matrix contaminated with a mixture of VOCs. *Environ Sci Technol* 46(11):5756–5763
- Liu Y, Whitman WB (2008) Metabolic, phylogenetic and ecological diversity of the methanogenic archaea. *Ann NY Acad Sci* 1125(1):171–189
- Liu A, Garcia-Dominguez E, Rhine ED, Young LY (2004) A novel arsenate respiring isolate that can utilize aromatic substrates. *FEMS Microbiol Ecol* 48(3):323–332
- Lovley DR, Giovannoni SJ, White DC, Champine JE, Phillips EJP, Gorby YA, Goodwin S (1993) *Geobacter metallireducens* gen. nov. sp. nov., a microorganism capable of coupling the complete oxidation of organic compounds to the reduction of iron and other metals. *Arch Microbiol* 159(4):336–344
- Morris BEL, Henneberger R, Huber H, Moissl-Eichinger C (2013) Microbial syntrophy: interaction for the common good. *FEMS Microbiol Rev* 37(3):384–406
- Nazina TN, Tourova TP, Poltarau AB, Novikova EV, Grigoryan AA, Ivanova AE, Lysenko AM, Petrunyaka VV, Osipov GA, Belyaev SS, Ivanov MV (2001) Taxonomic study of aerobic thermophilic bacilli: descriptions of *Geobacillus subterraneus* gen. nov., sp. nov. and *Geobacillus uzenensis* sp. nov. from petroleum reservoirs and transfer of *Bacillus stearothermophilus*, *Bacillus thermo-catenulatus*, *Bacillus thermoleovorans* and *Bacillus kaustophilus*, *Bacillus thermoglucosidasius* and *Bacillus thermodenitrificans* to *Geobacillus* as the new combinations *G. stearothermophilus*, *G. thermocatenulatus*, *G. thermoleovorans*, *G. kaustophilus*, *G. thermoglucosidasius* and *G. thermodenitrificans*. *Int J Syst Evol Microbiol* 51(2):433–446
- Nevin KP, Lovley DR (2002) Mechanisms for accessing insoluble Fe(III) oxide during dissimilatory Fe(III) reduction by *Geothrix fermentans*. *Appl Environ Microbiol* 68(5):2294–2299
- Nossa CW, Obedorf WE, Yang L, AAS JA, Paster BJ, Desantis TZ, Brodie EL, Malamud D, Poles MA, Pei Z (2010) Design of 16S rRNA gene primers for 454 pyrosequencing of the human foregut microbiome. *World J Gastroenterol* 16(33):4135–4144
- O'Sullivan LA, Mahenthalingam E (2005) Biotechnological potential within the genus *Burkholderia*. *Lett Appl Microbiol* 41(1):8–11
- Patureau D, Godon JJ, Bouchez T, Bernet N, Delgenes JP, Moletta R (1998) *Microvirgula aerodenitrificans* gen. nov.,

- sp. nov., a new gram-negative bacterium exhibiting co-respiration of oxygen and nitrogen oxides up to oxygen-saturated conditions. *Int J Syst Evol Microbiol* 48(3): 775–782
- Philippe G, Vega D, Bastide J (2001) Microbial hydrolysis of methyl aromatic esters by *Burkholderia cepacia* isolated from soil. *FEMS Microbiol Ecol* 37(3):251–258
- Rakoczy J, Schleinitz KM, Müller N, Richnow HH, Vogt C (2011) Effects of hydrogen and acetate on benzene mineralization under sulphate-reducing conditions. *FEMS Microbiol Ecol* 77(2):238–247
- Ramamoorthy S, Sass H, Langner H, Schumann P, Kroppenstedt RM, Spring S, Overmann J, Rosenzweig RF (2006) *Desulfosporosinus lacus* sp. nov., a sulfate-reducing bacterium isolated from pristine freshwater lake sediments. *Int J Syst Evol Microbiol* 56(12):2729–2736
- Ramos DT, Da Silva MLB, Chiaranda HS, Alvarez PJJ, Corseuil HX (2013) Biostimulation of anaerobic BTEX biodegradation under fermentative methanogenic conditions at source-zone groundwater contaminated with a biodiesel blend (B20). *Biodegradation* 24(3):333–341
- Robertson WJ, Bowman JP, Franzmann PD, Mee BJ (2001) *Desulfosporosinus merdicii* sp. nov., a spore-forming sulfate-reducing bacterium isolated from gasoline-contaminated groundwater. *Int J Syst Evol Microbiol* 51:133–140
- Sanford RA, Cole JR, Tiedje JM (2002) Characterization and description of *Anaeromyxobacter dehalogenans* gen. nov., sp. nov., an aryl-halo-respiring facultative anaerobic myxobacterium. *Appl Environ Microbiol* 68(2):893–900
- Schink B (1997) Energetics of syntrophic cooperation in methanogenic degradation. *Microbiol Mol Biol Rev* 61(2):262–280
- Shim H, Hwang B, Lee S, Kong S (2005) Kinetics of BTEX biodegradation by a coculture of *Pseudomonas putida* and *Pseudomonas fluorescens* under hypoxic conditions. *Biodegradation* 16(4):319–327
- Slobodkin AI (2005) Thermophilic microbial metal reduction. *Microbiology* 74(5):501–504
- Smith KS, Ingram-Smith C (2007) *Methanosaeta*, the forgotten methanogen? *Trends Microbiol* 15(4):150–155
- Sousa DZM, Pereira AM, Stams AJM, Alves MM, Smidt H (2007) Microbial communities involved in anaerobic degradation of unsaturated or saturated long-chain fatty acids. *Appl Environ Microbiol* 73(4):1054–1064
- Sousa DZM, Smidt H, Alves MM, Stams AJM (2009) Ecophysiology of syntrophic communities that degrade saturated and unsaturated long-chain fatty acids. *FEMS Microbiol Ecol* 68(3):257–272
- Spalding BP, Watson DB (2006) Measurement of dissolved H₂, O₂ and CO₂ in groundwater using passive samplers for gas chromatographic analyses. *Environ Sci Technol* 40(24): 7861–7867
- Spalding BP, Watson DB (2008) Passive sampling and analyses of common dissolved fixed gases in groundwater. *Environ Sci Technol* 42(10):3766–3772
- Tanaka K, Stackebrandt E, Tohyama S, Eguchi T (2000) *Desulfovira adipica* gen. nov., sp. nov., an adipate-degrading, gram-negative, sulfate-reducing bacterium. *Int J Syst Evol Microbiol* 50(2):639–644
- Thauer RK, Jungermann K, Decker K (1977) Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol Rev* 41(1):100–180
- Treude N, Rosencrantz D, Liesack W, Schnell S (2003) Strain FAc12, a dissimilatory iron-reducing member of the Anaeromyxobacter subgroup of *Myxococcales*. *FEMS Microbiol Ecol* 44(2):261–269
- van der Zaan BM, Saia FT, Stams AJM, Plugge CM, de Vos WM, Smidt H, Langenhoff AAM, Gerritse J (2012) Anaerobic benzene degradation under denitrifying conditions: *Peptococcaceae* as dominant benzene degraders and evidence for a syntrophic process. *Environ Microbiol* 14(5):1171–1181
- Villemur R, Lanthier M, Beaudet R, Lépine F (2006) The *Desulfuobacterium* genus. *FEMS Microbiol Rev* 30(5):706–733
- Wagner ID, Wiegel J (2008) Diversity of thermophilic anaerobes. *Ann NY Acad Sci* 1125(1):1–43
- Ward NL, Challacombe JF, Janssen PH, Henrissat B, Coutinho PM, Wu M, Xie G, Haft DH, Sait M, Badger J, Barabote RD, Bradley B, Brettin TS, Brinkac LM, Bruce D, Creasy T, Daugherty TM, Deboy RT, Dettler JC, Dodson RJ, Durkin AS, Ganapathy A, Dwinin-Giglio M, Han CS, Khouri H, Kiss H, Kothari SP, Madupu R, Nelson KE, Nelson WC, Paulsen I, Penn K, Ren Q, Rosovitz MJ, Selengut JD, Shrivastava S, Sullivan SA, Tapia R, Thompson LS, Watkins KL, Yang Q, Yu C, Zafar N, Zhou L, Kuske CR (2009) Three genomes from the phylum Acidobacteria provide insight into lifestyles of these microorganisms in soils. *Appl Environ Microbiol* 75(7):2046–2056
- Worm P, Müller N, Plugge CM, Stams AJM, Schink B (2011) Syntrophy in methanogenic degradation. *Microbiol Monogr* 19:143–173
- Zhang X, Peterson C, Reece D, Moller G, Haws R (1998) Biodegradability of biodiesel in the aquatic environment. *Trans Am Soc Agric Eng* 41:1423–1430